

AD-A138 895

THE POSTRESUSCITATION PATIENT: VENTILATION AND  
METABOLISM(U) COLUMBIA UNIV NEW YORK DEPT OF SURGERY  
J M KINNEY OCT 81 DAMD17-81-C-1048


1/1

UNCLASSIFIED

F/G 6/16

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A138895

PROGRESS REPORT

Contract # DAMD17-81-C-1048

October 1, 1980 - September 30, 1981

U.S. ARMY RESEARCH & DEVELOPMENT COMMAND

John M. Kinney, M.D.  
Principal Investigator

Department of Surgery  
College of Physicians and Surgeons  
Columbia University  
New York City

Approved for public release; distribution unlimited

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

DTIC FILE COPY

DTIC  
ELECTE  
S  
MAR 12 1984  
A

84 03 09 076

## PROGRESS REPORT

Contract # DAMD17-81-C-1048

October 1, 1980 - September 30, 1981

U.S. ARMY RESEARCH & DEVELOPMENT COMMAND

John M. Kinney, M.D.  
Principal Investigator

Department of Surgery  
College of Physicians and Surgeons  
Columbia University  
New York City

Approved for public release; distribution unlimited

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.



Application For

SWAMI

TAN

1940

1941

1942

1943

1944

1945

1946

1947

1948

1949

1950

1951

1952

1953

1954

1955

1956

1957

1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971

1972

1973

1974

1975

1976

1977

1978

1979

1980

1981

1982

1983

1984

1985

1986

1987

1988

1989

1990

1991

1992

1993

1994

1995

1996

1997

1998

1999

2000

2001

2002

2003

2004

2005

2006

2007

2008

2009

2010

2011

2012

2013

2014

2015

2016

2017

2018

2019

2020

2021

2022

2023

2024

2025

2026

2027

2028

2029

2030

2031

2032

2033

2034

2035

2036

2037

2038

2039

2040

2041

2042

2043

2044

2045

2046

2047

2048

2049

2050

2051

2052

2053

2054

2055

2056

2057

2058

2059

2060

2061

2062

2063

2064

2065

2066

2067

2068

2069

2070

2071

2072

2073

2074

2075

2076

2077

2078

2079

2080

2081

2082

2083

2084

2085

2086

2087

2088

2089

2090

2091

2092

2093

2094

2095

2096

2097

2098

2099

2100

2101

2102

2103

2104

2105

2106

2107

2108

2109

2110

2111

2112

2113

2114

2115

2116

2117

2118

2119

2120

2121

2122

2123

2124

2125

2126

2127

2128

2129

2130

2131

2132

2133

2134

2135

2136

2137

2138

2139

2140

2141

2142

2143

2144

2145

2146

2147

2148

2149

2150

2151

2152

2153

2154

2155

2156

2157

2158

2159

2160

2161

2162

2163

2164

2165

2166

2167

2168

2169

2170

2171

2172

2173

2174

2175

2176

2177

2178

2179

2180

2181

2182

2183

2184

2185

2186

2187

2188

2189

2190

2191

2192

2193

2194

2195

2196

2197

2198

2199

2200

2201

2202

2203

2204

2205

2206

2207

2208

2209

2210

2211

2212

2213

2214

2215

2216

2217

2218

2219

2220

2221

2222

2223

2224

2225

2226

2227

2228

2229

2230

2231

2232

2233

2234

2235

2236

2237

2238

2239

2240

2241

2242

2243

2244

2245

2246

2247

2248

2249

2250

2251

2252

2253

2254

2255

2256

2257

2258

2259

2260

2261

2262

2263

2264

2265

2266

2267

2268

2269

2270

2271

2272

2273

2274

2275

2276

2277

2278

2279

2280

2281

2282

2283

2284

2285

2286

2287

2288

2289

2290

2291

2292

2293

2294

2295

2296

2297

2298

2299

2300

2301

2302

2303

2304

2305

2306

2307

2308

2309

2310

2311

2312

2313

2314

2315

2316

2317

2318

2319

2320

2321

2322

2323

2324

2325

2326

2327

2328

2329

2330

2331

2332

2333

2334

2335

2336

2337

2338

2339

2340

2341

2342

2343

2344

2345

2346

2347

2348

2349

2350

2351

2352

2353

2354

2355

2356

2357

2358

2359

2360

2361

2362

2363

2364

2365

2366

2367

2368

2369

2370

2371

2372

2373

2374

2375

2376

2377

2378

2379

2380

2381

2382

2383

2384

2385

2386

2387

2388

2389

2390

2391

2392

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
AD-A138895		
4. TITLE (and Subtitle) THE POSTRESUSCITATION PATIENT: VENTILATION AND METABOLISM		5. TYPE OF REPORT & PERIOD COVERED Annual Progress Report October 1, 1980 - September 30, 1981
7. AUTHOR(s) John M. Kinney, M.D.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Surgery College of Physicians and Surgeons Columbia University, New York City, New York		8. CONTRACT OR GRANT NUMBER(s) DAMD17-81-C-1048
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command Fort Detrick Frederick, Maryland 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62772A.3S162772A874.AA.125
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE October 1981
		13. NUMBER OF PAGES 26
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		

PROGRESS REPORT

TABLE OF CONTENTS

PAGE

I.	Publications	5
II.	Effects of Nutrition and Injury on Ventilation	8
III.	Muscle Composition	20
IV.	References	26

## I PUBLICATIONS

### Journals

1. Askanazi J, Rosenbaum SH, Michelsen CB, Elwyn DH, Hyman AI, Kinney JM:  
Increased body temperature secondary to total parenteral nutrition.  
Crit Care Med 8:736-737, 1980
2. Carpentier YA, Askanazi J, Elwyn DH, Gump FE, Nordenstrom J, Kinney JM:  
The effect of carbohydrate intake on the lipolytic rate in depleted  
patients. Metabolism 29:974-979, 1980
3. Robin AP, Nordenstrom J, Askanazi J, Elwyn DH, Carpentier YA, Kinney JM:  
Plasma clearance of fat emulsion in trauma and sepsis: Use of a  
three-stage lipid clearance test. JPEN 4:505-510, 1980
4. Elwyn DH: Nutritional requirements of adult surgical patients. Crit  
Care Med 8:9-20, 1980
5. Askanazi J, Carpentier YA, Michelsen CB, Elwyn DH, Furst P, Kantrowitz LR,  
Gump FE, Kinney JM: Muscle and plasma amino acids following injury:  
Influence of intercurrent infection. Ann Surg 192:78-85, 1980
6. Askanazi J, Carpentier YA, Jeevanandam M, Michelsen CB, Elwyn DH, Kinney JM:  
Energy expenditure, nitrogen balance, and norepinephrine excretion  
after injury. Surgery 89:478-484, 1981
7. Askanazi J, Nordenstrom J, Rosenbaum SH, Elwyn DH, Hyman AI, Carpentier YA,  
Kinney JM: Nutrition for the patient with respiratory failure:  
Glucose vs fat. Anesthesiology 54:373-377, 1981

8. Bergstrom J, Furst P, Holmstrom B, Vinnars E, Askanazi J, Elwyn DH, Michelsen CB, Kinney JM: Influence of injury and nutrition on muscle water and electrolytes: Effect of elective operation. Ann Surg 193:810-816, 1981
9. Carpentier YA, Nordenstrom J, Robin AP, Kinney JM: Glycerol turnover and kinetics of exogenous fat in surgical patients, in Wright PD (ed): Acta Chir Suppl 507, Symposia, 2nd European Congress on Parenteral and Enteral Nutrition, Newcastle upon Tyne, September 7-10, 1980, Stockholm, Almqvist & Wiksell Periodical Company, 1981, p 226
10. Robin AP, Carpentier YA, Askanazi J, Kinney JM: Metabolic consequences of hypercaloric glucose infusions: A review. Acta Chir Belg 80, 1981
11. Robin AP, Askanazi J, Greenwood MRC, Carpentier YA, Gump FE, Kinney JM: Lipoprotein lipase activity in surgical patients: Influence of trauma and infection. Surgery 80:401-408, 1981
12. Robin AP, Askanazi J, Cooperman A, Carpentier YA, Elwyn DH, Kinney JM: Influence of hypercaloric glucose infusions on fuel economy in surgical patients: A review. Crit Care Med 9:680-686, 1981
13. Elwyn DH, Kinney JM, Gump FE, Jeevanandam M, Chikenji T, Askanazi J: Metabolic and endocrine effects of fasting followed by infusion of five percent glucose. Surgery (in press)
14. Nordenstrom J, Jeevanandam M, Elwyn DH, Carpentier YA, Askanazi J, Robin AP, Kinney JM: Increasing glucose intake during total parenteral nutrition increases norepinephrine excretion in trauma and sepsis. Clin Physiol (in press)
15. Liaw KY, Askanazi J, Michelsen CB, Furst PF, Elwyn DH, Kinney JM: Effect of postoperative nutrition on muscle high energy phosphates. Ann Surg (in press)



Chapters

1. Elwyn DH, Kinney JM, Askanazi J: Energy expenditure in surgical patients, in Mullen JL (ed): Surgical Clinics of North America, Vol 61, Philadelphia, WB Saunders Co, 1981, p 545
2. Furst P, Bergstrom J, Hellstrom B, Vinnars E, Herfarth C, Klippel C, Merkel N, Schultis K, Elwyn DH, Hardy M, Kinney JM: Amino acid metabolism in cancer, in Kluthe R, Lohr G-W (eds): Nutrition and Metabolism in Cancer, International Workshop, Freiburg 1979, Stuttgart - New York, Georg Thieme Verlag, 1981, p 75
3. Kinney JM, Elwyn DH: Protein-energy interrelationships, in Selvey N, White PL (eds): Nutrition in the 1980's: Constraints on our Knowledge. Proceedings of the Western Hemisphere Nutrition Congress VI, Los Angeles, August 10-14, 1980, New York, Alan R Liss Inc, 1981, p 179

## II EFFECTS OF NUTRITION AND INJURY ON VENTILATION.

A. Ventilatory studies in injured and depleted patients. Our previous studies with total parenteral nutrition have largely been based on a mixture of hypotonic glucose and amino acids which our group has shown to cause a significant increase in  $\text{CO}_2$  output. An associated increase in oxygen consumption indicating increased energy expenditure, does not occur in patients with nutritional depletion but does occur, superimposed on whatever level of hypermetabolism is present, if the patient is acutely injured or has a serious infection. Last year, we proposed to examine this problem by studying three groups of patients: normal convalescents following colon resection; normal convalescents following total hip replacement; and patients with major accidental injury. It was planned that within 48 hours following operation or injury the patients would be randomly assigned to receive 5% dextrose 20% dextrose or 20% dextrose plus amino acids. Various ventilatory measurements would be made at 6, 24 and 48 hours after starting the intravenous nutrition.

During January of 1981, these studies were begun, after completing a study of 6 depleted surgical patients whose ventilatory response to  $\text{CO}_2$  was studied while receiving 5% dextrose and again after 7 days of either of two hypercaloric intravenous diets providing energy equal to 1.5 times the REE and either 12 or 24 gms of amino acid nitrogen per day. The order of administration of the high and low nitrogen diets were randomized. Typical results are plotted for 6 patients in Figure 1. During the 5% dextrose infusions, these patients were less sensitive to  $\text{CO}_2$  than normal subjects. The effect of the low nitrogen total parenteral nutrition (TPN) was to increase the sensitivity of the patient to  $\text{CO}_2$ , an effect which was more marked




Figure 1

after 7 days of the high nitrogen intake.

Studies of acutely injured patients were begun with the administration of TPN to a young healthy male 3 days after receiving multiple gunshot wounds to the abdomen. He had extensive peritoneal contamination from multiple bullet wounds to the intestinal tract which were treated at emergency laparotomy with primary closure. He ran a low grade fever during the first postoperative week but then had a successful convalescence. Because of the probability of intraperitoneal sepsis, and prolonged delay in the use of his intestinal tract, he was started on TPN on his third postinjury day. His minute ventilation showed a dramatic response to TPN. On 5 % dextrose it was 8 and 10 l/min on room air and 2% CO<sub>2</sub> respectively. On TPN these values doubled to 16 and 20 l/min.

B. Effects of mouth piece and noseclip and of postural position on the respiratory response to CO<sub>2</sub>. It seemed to us that our studies on depleted patients and our studies with this gunshot victim indicated that carbohydrate infusion was stimulating hyperventilation mainly by increasing CO<sub>2</sub> production, while administration of amino acids was stimulating ventilation by an entirely different mechanism, perhaps by increasing the sensitivity of the ventilatory center to CO<sub>2</sub>. These studies were reviewed with Dr. Joseph Milic-Emili, Professor of Physiology at McGill University and an international authority on the control of breathing. Dr. Milic-Emili found the studies to be extremely provocative but emphasized that we needed further validation of our non-invasive system for measurement of gas exchange for use in spirometric studies. Our previous published validation was only with resting normal subjects and did not include CO<sub>2</sub> response curves using conventional equipment (mouthpiece and noseclip) in the same individuals who were studied in the canopy. He also thought that we must show the different effects of the supine and sitting position on respiration in normal man, in order for our data to allow quick comparison with studies in the literature. He also reviewed our plans for the years work on ventilation and metabolism and observed that our experimental design would allow no definition of the effect of amino acids given alone. Therefore, studies were undertaken to examine the influence of the mouthpiece and noseclip compared with the canopy in studying the CO<sub>2</sub> response in normal subjects and patients. A study of 8 supine male subjects and 6 patients has been completed during this contract year.

In random order, 8 supine normal subjects and 6 patients convalescing from major abdominal operations, with mean body surface areas of 1.88 and 1.70 respectively were studied in the canopy with a (a) mouthpiece plus

noseclip and (b) without any attachment to the face or airway. Ventilatory patterns were assessed during room air breathing and during the administration of 2% and 4% CO<sub>2</sub>. Measurements were made over a 10 minute period once stable levels of oxygen consumption and carbon dioxide production had been achieved. Ventilation ( $\dot{V}_E$ ), tidal volume ( $V_T$ ), frequency (f), inspiratory and expiratory times ( $T_I, T_E$ ) and peak inspiratory and expiratory flows were measured. Mean inspiratory and expiratory flows ( $V_T/T_I, V_T/T_E$ ) and inspiratory duty cycle ( $T_I/T_{TOT}$ ) were calculated.

Normal subjects showed a relatively predictable increase in  $\dot{V}_E$ ,  $V_T$  and  $V_T/T_I$  with the use of a MP + NC (Table I). Patients showed a more variable and unpredictable response to a mouthpiece plus noseclip.

Table I

NORMALS (n = 8)

		$\dot{V}_E$ (ml/min)	$V_T$ (ml)	f (min <sup>-1</sup> )	$V_T/T_I$ (ml/sec)	$T_I/T_{TOT}$
CANOPY	RA	5.6	379	16	254	.38
	2% CO <sub>2</sub>	7.0	471	16	304	.39
	4% CO <sub>2</sub>	12.3	716	17	511	.41
MP+NC	RA	6.7*	444*	17	292*	.38
	2% CO <sub>2</sub>	8.0*	488	17	317*	.40
	4% CO <sub>2</sub>	13.8	761	18	559	.42

PATIENTS (n = 6)

CANOPY	RA	6.5	277	23	257	.39
	2% CO <sub>2</sub>	8.9	383	22	322	.41
	4% CO <sub>2</sub>	13.1	593	22	504	.42
MP+NC	RA	6.0	271	21	293	.41
	2% CO <sub>2</sub>	8.5	396	21	343	.41
	4% CO <sub>2</sub>	12.1	548	22	482	.42

Values are means; \* significantly different from measurements with canopy alone (p<.05).

Previous studies have shown that a mouthpiece plus noseclip increased  $V_T$  16-29%, and  $\dot{V}_E$  14-31% in normal subjects breathing room air. In the present study ventilatory apparatus induced an increase in minute ventilation during both room air and stimulated breathing in normal subjects. These increases in ventilation are due to increases in tidal volume secondary to increases in mean inspiratory flow. In the surgical patients, the mouthpiece plus noseclip failed to induce any consistent changes in breathing patterns.

The surgical patients had higher frequencies and lower tidal volumes at rest, as has been previously reported. The response to  $CO_2$  as measured by changes in  $\dot{V}_E$  was similar in patients and normals. While breathing  $CO_2$  the characteristic changes induced by a mouthpiece plus noseclip were observed in normal subjects but not in patients. The changes in the control of breathing which led to a high  $f$  and decreased  $V_T$  in the patients may have attenuated the stimulus of the mouthpiece plus noseclip. Thus, the MP+NC had a predictable effect on breathing in normal subjects, while in patients it had a variable and less predictable effect. This data has important implications for studies of control of breathing in both normal subjects and surgical patients.

In order to meet the other question of Dr. Milic-Emili, regarding validation of our equipment, nine normal subjects have been studied in both the supine and seated position while breathing room air and during the administration of 2 and 4%  $CO_2$ . The minute ventilation was 21% higher in the sitting position, and the tidal volume and mean inspiratory flow rates were also greater. When the relationship between minute ventilation and mean inspiratory flow was examined the ratio was always higher in the

seated position. No difference in response to  $\text{CO}_2$  was seen when minute ventilation was compared with arterial  $\text{PCO}_2$  or when mean inspiratory flow was compared with arterial  $\text{PCO}_2$ . However, the arterial  $\text{PCO}_2$  in the supine position was significantly higher than in the seated position for every level of ventilation.

C. Effect of dextrose and amino acids on ventilatory chemosensitivity.

Our studies aimed at understanding the influence of intravenous glucose and intravenous amino acids when administered separately to normal and hyper-metabolic surgical patients have made good progress during this year.

In order to isolate the influence of glucose from the influence of amino acids in total parenteral nutrition, 4 healthy volunteer subjects were hospitalized and given intravenous glucose at either  $281 \text{ kcal/m}^2/24\text{hr}$  or  $1125 \text{ kcal/m}^2/24\text{hr}$  for 3 days each in random order. On the third day of each diet, ventilation and gas exchange were measured with room air, and following the administration of 2 and 4%  $\text{CO}_2$ . Arterial blood gases were sampled during each measurement. Ventilatory chemosensitivity to  $\text{CO}_2$  was calculated as  $\Delta V_E / \Delta \text{PaCO}_2$  during  $\text{CO}_2$  inhalation. High glucose infusion, while breathing room air, caused increases of 23% in minute ventilation, 27% in tidal volume, 12% in inspiratory flow rate and 13% in  $\text{CO}_2$  production, as compared to values obtained with the low glucose infusion rate. Oxygen consumption and respiratory frequency were unchanged between the two levels of glucose infusion and there was no difference in ventilatory chemosensitivity. Therefore, in normal subjects, the increase in minute ventilation associated with glucose intake appears to be secondary to the increase in  $\text{CO}_2$  (due to increasing RQ) but no change occurs in the behavior of the respiratory center to this increase in  $\text{CO}_2$  production. To further

investigate the effect of nutrients on ventilation and examine the correlation between resting energy expenditure and chemosensitivity the effects of isotonic amino acid (AA) infusions by peripheral vein on  $O_2$  consumption,  $CO_2$  production and the ventilatory response to  $CO_2$  were studied. The effects of a four hour amino acid infusion were studied in convalescing surgical patients and the data compared to that observed in normal subjects.

Six normal subjects (Group I) and five patients (Group II) convalescing from major surgery were studied. In group I,  $CO_2$  responses were assessed after an overnight fast (control) and following 4 hours of an infusion of 125 ml/hr of an isotonic amino acid solution (AA<sub>4</sub>), via peripheral vein (Aminosyn 3.5%).

The patients all had infusions of 5% dextrose for 24 hours or more followed by identical infusions of isotonic AA for 4 hours (AA<sub>4</sub>). The response to  $CO_2$  was evaluated prior to and after 4 hours of amino acids.

The normal subjects had a 54% ( $p < .05$ ) increase in ventilatory sensitivity as measured by the change in  $VE$ /change in  $PaCO_2$  during  $CO_2$  administration. This was associated with an 11% increase in resting energy expenditure (REE). The patients had an 84% increase in chemosensitivity with a 2% increase in REE (Table II).

Isotonic amino acids administered in the postabsorptive state to normal subjects increase sensitivity to  $CO_2$  within 4 hours. This effect does not appear to be due to changes in pH, but is associated to changes in metabolic rate in normals. The patients had a proportionally greater increase in ventilatory sensitivity than energy expenditure with the amino acid infusions. This indicates that an increase in metabolic rate may not entirely explain the phenomenon.



Table II

GROUP I

Normal Subjects (n=6)

	$\frac{\dot{V}_E}{PaCO_2}$ (1/torr- CO <sub>2</sub> )	$\frac{V_T/TI}{PaCO_2}$ (ml/sec torr CO <sub>2</sub> )	REE (kcal/24* hour)	HCO <sub>3</sub> $\frac{meq}{l}$
Control	2.2 ± 1.2	65 ± 56	1302 ± 410	26 ± 2
AA <sub>4</sub>	3.4 ± 2.0	110 ± 60	1441 ± 399	25 ± 2

GROUP II

Patients (n=5)

	$\frac{\dot{V}_E}{PaCO_2}$ (1/torr- CO <sub>2</sub> )	$\frac{V_T/TI}{PaCO_2}$ (ml/sec torr CO <sub>2</sub> )	REE (kcal/24* hour)	HCO <sub>3</sub> $\frac{meq}{l}$
Control	3.2 ± 1.2	131 ± 100	1501 ± 283	27 ± 3
AA <sub>4</sub>	6.0 ± 4.9	230 ± 190	1536 ± 210	27 ± 2

all values are mean ± SD

\*calculated from V<sub>O</sub><sub>2</sub>, V<sub>CO</sub><sub>2</sub> and RQ

These results suggest that peripheral infusions of amino acids enhance ventilatory drive as measured by the response to CO<sub>2</sub> after four hours of infusion. This data has important implications in the care of the patient with CO<sub>2</sub> retention. Since the amino acid infusions are administered by peripheral vein this phenomenon may have a potential for broad application.

Our application to the Army last year referred to the data in the literature suggesting that normal subjects receiving only 5% dextrose by vein in amounts of approximately 100 gms per day will develop decreased ventilatory responses to hypoxia and perhaps to hypercarbia within 7 to 10 days. This year we have undertaken a study of 8 normal subjects with the dual objectives of confirming this finding and also of determining how rapidly an amino acid infusion might act to return the CO<sub>2</sub> response to normal, if it is indeed reduced. We have hospitalized 8 normal subjects who




Figure 3




Figure 2

received approximately 100 grams of glucose a day as their sole nutrition for 1 week. At the end of that time they received an amino acid infusion for 24 hours. Gas exchange and ventilatory studies including the response to  $\text{CO}_2$  were performed on Day 0, and Day 7 as well as after 24 hours of amino acid infusion. The subjects showed a distinct decrease in their resting energy expenditure during the week on 5% dextrose with a corresponding decrease in minute ventilation and tidal volumes (Figs. 2 & 3). Amino acid infusion caused a return toward control values within 4 hours which was complete within 24 hours.

D. Use and calibration of commercial respiratory equipment. Measurements of  $\text{O}_2$  consumption ( $\dot{V}\text{O}_2$ ),  $\text{CO}_2$  production ( $\dot{V}\text{CO}_2$ ) and respiratory quotient (RQ) have been demonstrated to be important clinical parameters, particularly in patients with respiratory failure. Performing these measurements in patients receiving mechanical ventilation has proved difficult. Recently, two commercial instruments (Siemens-Elema Servo Ventilator 900B and Beckman Metabolic Measurement Cart) designed to perform measurements of gas exchange during mechanical ventilation have been introduced. However, the accuracy and reproducibility of these instruments remains to be evaluated. We examined the capabilities of these instruments and devised a simple model for initial testing and continued recalibration of instruments designed to perform gas exchange measurements.

In the initial phase both instruments were used to ventilate a lung model (Fig. 4). Gas exchange was reproduced by administration of  $\text{CO}_2$  (to simulate  $\dot{V}\text{CO}_2$ ) and  $\text{N}_2$  (to simulate  $\dot{V}\text{O}_2$ ). Flows of both gases were introduced into the system using Matheson flowmeters (601 for  $\text{CO}_2$  and 603 for  $\text{N}_2$ ) which were calibrated using a mercury-O-ring dry spirometer. Ventilatory conditions were varied (frequency = 10, 15, 10, tidal volume =




Figure 5

Figure 4

250, 333, 350, 466, 500, 666, 700 and 1000 ml,  $F_{I}O_2 = .21, .40, .85$ ) as were the simulated levels of gas exchange ( $V_{CO_2} = 85 - 270$  ml/min,  $V_{O_2} = 80 - 933$  ml/min). Each condition was studied for 8 minute periods on 4 occasions. In the second phase of the study the ability of the two instruments to measure RQ was assessed by burning methanol ( $RQ = .67$ ) and ventilating the lung model as shown (Fig. 5).

Representative results using ventilatory parameters and levels of gas exchange that would be expected in an adult patient are shown in Fig. 6. At physiological tidal volumes for adults (466-700 ml) there was less than a 7% difference between measured vs administered  $CO_2$ . With smaller tidal volumes (150-350 ml) the discrepancies widened to 13% and 10% respectively for the Siemens (S/E) and Beckman (B) systems respectively. Measured vs calculated  $O_2$  consumption was within 10% and 5% for the S/E and B systems on room air. With increasing levels of  $F_{I}O_2$  the difference between measured and known RQ remained within 5% and 1.5% for S/E and B respectively.




Figure 6

Thus both instruments give accurate and reproducible measurements of gas exchange as assessed by this in vitro testing system. This system appears to be useful for initial testing and continued validation of these instruments.

It is important to emphasize that both the S/E Servo Ventilator and the B Metabolic Measurement Cart are precision instruments that require careful calibration and continued maintenance. Both instruments, even when calibrated according to the specifications of the manufacturer, can periodically yield unreliable data. Thus a routine system for continued validation is mandatory.

### III Muscle Composition.

A. Muscle water, electrolytes and glycogen. Prior studies from this unit (see progress report 1979-1980) have shown that muscle water, chloride and sodium increase after severe injury or during sepsis. This indicates an expansion of the extracellular compartment of muscle similar to that measured by whole body tracer studies.(1). These data, combined with data of Bergström, Fürst and Vinnars, collected in Sweden, are currently being prepared for publication. Other studies from this laboratory (2,3, Progress Report 1979-1980) have shown an interrelation between the effects of diet and injury on high energy phosphate compounds in muscle. Fasting normal subjects produces no change. Fasted patients, after total hip operation, have reduced concentrations of ATP and ADP and increased concentrations of AMP, however these changes are abolished by infusing 5% dextrose. Severely injured or septic patients show even bigger changes in high energy phosphates which are not abolished by 5% dextrose but are overcome with total parenteral nutrition. Thus with injury there are decreases in ATP and ADP and increases in AMP which are more marked with increasing severity. Glucose administration can

Table III. Muscle water electrolytes. Membrane potential set at 87.2 mv.

	Normal	Low Glucose	High Glucose
n	37	7	7
Water - ml/100 g FFS			
Total	333.4 $\pm$ 1.9	365.2 $\pm$ 12.1***	345.4 $\pm$ 11.4
Extracellular	45.7 $\pm$ 1.5	103.1 $\pm$ 20.2***	87.0 $\pm$ 18.8***
Intracellular	287.7 $\pm$ 2.4	262.1 $\pm$ 19.3**	258.5 $\pm$ 15.3**
Electrolytes - meq/100 g FFS			
Sodium	9.9 $\pm$ .3	14.0 $\pm$ 2.3**	11.6 $\pm$ 1.1
Potassium	45.8 $\pm$ .2	45.7 $\pm$ .2	44.2 $\pm$ .9**
Magnesium	8.4 $\pm$ .1	8.6 $\pm$ .1	8.4 $\pm$ .2
Chloride	6.5 $\pm$ .2	12.2 $\pm$ 2.2***	10.2 $\pm$ 1.9***
Electrolytes - meq/l IC water			
Sodium	11.7 $\pm$ 1.1	.4 $\pm$ 5.7*	-.8 $\pm$ 6.2**
Potassium	158.9 $\pm$ 1.3	172.1 $\pm$ 13.6	171.0 $\pm$ 7.2**
Magnesium	29.1 $\pm$ .5	32.0 $\pm$ 2.5	32.3 $\pm$ 1.3*

Differs from normal,  $p < .05^*$ ,  $.01^{**}$ ,  $.001^{***}$

inhibit these changes, and is more effective as the amount increases. Since membrane potential is closely linked to ATP and ADP concentrations, and electrolyte and water distribution are in turn dependent on membrane potential, it seemed likely that diet and particularly glucose might affect the changes in water and electrolytes seen in injury and sepsis.

With these considerations in mind, a prospective study was initiated of the effects of low (550 kcal/day) and high (2100 kcal/day) glucose intakes on the changes in muscle water, electrolytes and glycogen induced by injury. Five severely injured patients (3 with gunshot wounds, 1 after a 5 story fall, and 1 postcystectomy) and 3 septic patients were maintained either for 3 days on the low intake followed by three days on the high intake, or vice versa. Muscle biopsies and blood samples were taken on the third day of each diet. Data for seven of the patients are given here. Compared to normal values these patients had significant increases for muscle water, sodium and chloride on the low glucose intake (Table III). On the high glucose diet the increases were smaller, only chloride showing a significant increase. There was a significant decrease in potassium on the high but not the low intake. These data suggest that high glucose may partially counteract the effects of injury, however with the small numbers used there was no significant differences between the high and low intakes by the paired t test.

The values for intracellular (ICW) and extracellular water (ECW) and for electrolyte concentrations in ICW have been calculated from the chloride data using the Nernst equation and assuming a normal membrane potential of 87.2 mv.(4) They indicate a marked expansion of ECW and contraction of ICW (Table III). However, it is unlikely that the membrane potential was not affected by the severe injury or sepsis. Furthermore the calculated decrease in IC sodium and increase in IC potassium concentrations are inconsistent with an unchanged membrane potential. It is not feasible to measure the membrane potential in these studies, however, various arbitrary values may



Table IV. Muscle water and electrolytes. Membrane potential set at 55 mv.

	Low Glucose	High Glucose
n	7	7
Water - ml/100 g FFS		
Total	365.2 $\pm$ 12.1	345.4 $\pm$ 11.4
Extracellular	75.3 $\pm$ 21.9	59.6 $\pm$ 20.0
Intracellular	289.9 $\pm$ 21.3	285.8 $\pm$ 16.9
Electrolytes - meq/l IC water		
Sodium	13.0 $\pm$ 5.1	12.0 $\pm$ 5.6
Potassium	156.1 $\pm$ 12.3	155.2 $\pm$ 6.5
Magnesium	29.1 $\pm$ 2.2	29.4 $\pm$ 1.2

be used in calculating ECW and ICW. When a membrane potential of 55 mv was used, IC sodium concentrations were increased and potassium concentrations were decreased from normal, qualitatively consistent with a reduced potential (Table IV). With this membrane potential, calculated values for ICW no longer differ from normal. However ECW values remain higher than normal and are higher on the low than the high glucose intake. It seems likely that, in these patients, there is both a change in membrane potential as well as an expansion of ECW.

Glycogen has been determined in muscle biopsies from the eight patients and from four normal subjects treated identically. In both groups there was a significant increase in glycogen content in going from the low to the high glucose intake (Table V). These changes are similar to those previously reported by Hultman. (5) The injured patients had lower muscle glycogen

Table V Muscle glycogen concentrations

	Low Glucose	High Glucose	Difference
	g glycogen/100 g fat free solids $\pm$ SEM		
Normals (n = 4)	4.7 $\pm$ 1.0	6.4 $\pm$ .6	1.7 $\pm$ .5*
Injured/septic (n = 8)	3.1 $\pm$ .7	5.9 $\pm$ .5	2.8 $\pm$ .3**

\*p<.05, \*\*p<.001

contents than the normals on the low diet and bigger increases in going to the high diet. However, these differences were of borderline significance and more studies are needed to confirm the findings.

We may conclude that increasing glucose intake increases glycogen content in injured/septic patients, possibly to a greater extent than in normal, and that this may be accompanied by partial prevention of the decreases in muscle membrane potential and increase in muscle ECW induced by injury or sepsis. More studies are planned for the coming year, both of normals and sick patients, to confirm these findings.

B. Muscle high energy phosphate compounds. During the past year the manual, enzymatic methods we were using for measurements of ATP, ADP, AMP, creatine, phosphocreatine, glucose-6-phosphate, and lactate in muscle biopsies have been adapted for use with a newly purchased Instrumentation Laboratories fast centrifugal analyzer. This permits more rapid and reproducible analyses on smaller amounts of material. No human studies were conducted during this period but we plan to resume them shortly.

C. Muscle concentrations of carnitine derivatives and of carnitine palmityl transferase. It has been suggested that part of the metabolic derangement seen in injury and sepsis may be due to an absolute or relative carnitine deficiency. Carnitine is required for transport of long chain fatty acids into mitochondria. A carnitine deficiency would lead to decreased ability to oxidize fat and increased reliance on protein and glucose for energy, and might help to explain the increased nitrogen losses in injured or septic patients. Border et al (6) have demonstrated that carnitine concentrations are reduced in sepsis in both man and dog.

With this in mind we have initiated a study of the effect of injury, sepsis and diet on muscle concentrations of free carnitine, carnitine esters, and the enzyme, carnitine palmityl transferase (CPT). During the past six months, methods for measuring these compounds (7,8,9) in open biopsies have been modified to permit accurate analysis in the small amount of tissue (ca 50 mg) which can be obtained with the noninvasive needle biopsy technique which we are using for measuring muscle water, electrolytes, glycogen, high energy phosphates and amino acids (10). These modifications have been completed and studies are now in progress on the effects of high and low glucose intakes in normal subjects and in injured or septic patients.

#### IV REFERENCES

1. Elwyn DH, Bryan-Brown CW, Shoemaker WC: Nutritional aspects of body water dislocation in postoperative and depleted patients. *Ann Surg* 182:76, 1975
2. Liaw KY, Askanazi J, Michelsen CB, et al: Effect of postoperative nutrition on muscle high energy phosphates. *Ann Surg* (in press)
3. Bergström J, Boström H, Fürst P, et al: Preliminary studies of energy-rich phosphagens in muscle from severely ill patients. *Crit Care Med* 4:197, 1976
4. Bergström J, Fürst P, Noree L-O, et al: Intracellular free amino acid concentration in human muscle tissue. *J Appl Physiol* 36:693, 1974
5. Hultman E: Muscle glycogen in man determined by needle biopsy specimen. *Scand J Clin Lab Invest* 19:209, 1967
6. Border JR, Burns GP, Rumph C, et al: Carnitine levels in severe infection and starvation: A possible key to the prolonged catabolic state. *Surgery* 68:175, 1970
7. Pearson DJ, Chase JFA, Tubbs PK, et al: The assay of (-)-carnitine and its O -acyl derivatives. *Methods Enzymol* 14:612, 1969
8. McGarry JD, Foster DW: An improved and simplified radioisotopic assay for the determination of free and esterified carnitine. *J Lipid Res.* 17:277, 1976
9. Nørum KR: Palmityl CoA: Carnitine palmityltransferase. Purification from calf-liver mitochondria and some properties of the enzyme. *Biochem Biophys Acta* 89:95, 1964
10. Bergström J: Chapter III obtaining the muscle samples. *Scand J Clin Lab Invest* 68:11, 1962

**FILMED**

**4-84**

**DTIC**